## Reported drug, alcohol, and cigarette use among high school students

By sex, race, ethnicity, and grade level, United States, 1995

(Percent reporting engaging in the behavior)

	Race, ethnicity										
	_	Sex			White, non- Black, non-			Grade level			
	Total	Male	Female	Hispanic	Hispanic	Hispanic	9th grade	10th grade	11th grade	12th grade	
Marijuana use, lifetime <sup>a</sup>	42.4%	45.2%	39.4%	40.5%	47.2%	49.2%	33.8%	41.4%	45.8%	47.0%	
Marijuana use, current <sup>b</sup>	25.3	28.4	22.0	24.6	28.6	27.8	20.9	25.5	27.6	26.2	
Cocaine use, lifetime <sup>c</sup>	7.0	8.8	5.0	6.5	2.0	16.0	5.7	7.5	7.2	7.4	
Cocaine use, current <sup>b</sup>	3.1	4.3	1.8	2.6	1.3	7.5	3.1	2.5	3.6	3.1	
Crack or freebase use, lifetime <sup>a</sup>	4.5	5.6	3.4	4.2	1.6	10.5	4.7	4.9	4.4	4.2	
Illegal steroid use, lifetime <sup>a</sup>	3.7	4.9	2.4	3.8	1.6	4.7	4.1	3.6	3.9	2.9	
Injected illegal-drug use, lifetime <sup>d</sup>	2.0	3.0	1.0	2.0	1.1	2.2	2.8	2.2	1.7	1.6	
Other illegal drug use, lifetime <sup>e</sup>	16.0	17.8	14.1	18.4	3.9	18.1	12.5	16.7	17.0	17.0	
Sniffed or inhaled intoxicating											
substances, lifetime <sup>f</sup>	20.3	22.1	18.4	22.7	9.5	22.8	24.6	22.4	19.2	15.9	
Alcohol use, lifetime <sup>g</sup>	80.4	81.1	79.5	81.7	73.7	82.9	72.4	78.9	83.2	85.5	
Alcohol use, currenth	51.6	53.2	49.9	54.1	42.0	54.7	45.6	49.5	53.7	56.5	
Episodic heavy drinking <sup>i</sup>	32.6	36.2	28.6	35.6	18.8	37.7	24.5	30.3	34.9	39.0	
Cigarette use, lifetime	71.3	72.1	70.4	71.1	66.0	76.3	63.4	71.1	75.8	73.8	
Cigarette use, currentk	34.8	35.4	34.3	38.3	19.2	34.0	31.2	33.1	35.8	38.2	
Cigarette use, frequent <sup>I</sup>	16.1	16.3	15.9	19.5	4.5	10.0	9.6	13.3	19.2	20.9	
Before age 13											
Smoked whole cigarette	24.9	27.8	21.8	25.9	17.2	26.6	28.1	25.6	25.0	21.5	
Drank alcohol <sup>m</sup>	32.4	38.6	25.5	30.3	35.9	39.5	41.0	34.1	29.6	25.8	
Tried marijuana	7.6	10.2	4.8	5.6	11.1	12.6	9.2	9.1	6.7	5.4	
Tried cocaine <sup>n</sup>	1.2	1.8	0.5	0.9	1.3	1.7	1.3	1.3	1.4	0.9	
On school property											
Cigarette use <sup>k</sup>	16.0	16.8	15.1	17.6	7.6	14.9	13.9	15.4	16.7	17.5	
Alcohol use <sup>K</sup>	6.3	7.2	5.3	5.6	7.6	9.7	7.5	6.0	5.7	6.2	
Marijuana use <sup>b</sup>	8.8	11.9	5.5	7.0	12.3	12.9	8.7	9.8	8.6	8.0	
Offered, sold, or given an illegal drug <sup>o</sup>	32.1	38.8	24.8	31.7	28.5	40.7	31.1	35.0	32.8	29.1	

Note: See Note, table 3.55. For survey methodology and sampling procedures, see Appendix 6.

Source: Laura Kann et al., "Youth Risk Behavior Surveillance--United States, 1995," CDC Surveillance Summaries, *Morbidity and Mortality Weekly Report* 45 No. SS-4 (Washington, DC: USGPO, Sept. 27, 1996), pp. 44-61. Table adapted by SOURCEBOOK staff.

<sup>&</sup>lt;sup>a</sup>Ever used. <sup>b</sup>Used one or more times during the 30 days preceding the survey.

<sup>&</sup>quot;Used one or more times during the su days preceding the survey."

Ever tried any form of cocaine (e.g., powder, crack, and freebase).

Ever injected illegal drugs. Respondents were classified as injecting-drug users only if they a) reported injecting-drug use not prescribed by a physician and b) answered "one or more" to any of these questions: "During your life, how many times have you used any form of cocaine including powder, crack, or freebase?"; "During your life, how many times have you used any other type of illegal drug, such as LSD, PCP, ecstacy, mushrooms, speed, ice, or heroin?"; "During your life, how many times have you taken steroid pills or shots without a doctor's prescription?"

<sup>&</sup>lt;sup>e</sup>Ever used any other type of illegal drug, such as LSD, PCP, ecstasy, mushrooms, speed,

<sup>&</sup>lt;sup>f</sup>Ever sniffed glue or breathed the contents of aerosol spray cans or inhaled any paint sprays to get high.

<sup>&</sup>lt;sup>g</sup>Ever had at least one drink of alcohol.

hDrank alcohol on 1 or more of the 30 days preceding the survey.

Drank five or more drinks of alcohol on at least one occasion on 1 or more of the 30 days preceding the survey.

Ever tried cigarette smoking, even one or two puffs.

On 1 or more of the 30 days preceding the survey.

Smoked cigarettes on 20 or more of the 30 days preceding the survey.

Other than a few sips.

On Including powder, crack, or freebase forms of cocaine.

ODuring the 12 months preceding the survey.